## §520.1860 Pradofloxacin.

- (a) Specifications. Each milliliter of suspension contains 25 milligrams (mg) pradofloxacin.
- (b) *Sponsor*. See No. 000859 in §510.600(c) of this chapter.
- (c) Special considerations. Federal law restricts this drug to use by or on the order of a licensed veterinarian. Federal law prohibits the extralabel use of this drug in food-producing animals.
- (d) Conditions of use in cats—(1) Amount. Administer 3.4 mg/lb (7.5 mg/kg) body weight once daily for 7 consecutive days.
- (2) Indications for use. For the treatment of skin infections (wounds and abscesses) in cats caused by susceptible strains of Pasteurella multocida, Streptococcus canis, Staphylococcus aureus, Staphylococcus felis, and Staphylococcus pseudintermedius.

[77 FR 76863, Dec. 31, 2012]

## §520.1870 Praziquantel tablets.

- (a) Specifications. Each tablet contains:
  - (1) 34 milligrams (mg) praziquantel.
  - (2) 11.5 or 23 mg praziquantel.
- (b) Sponsors. See sponsors in §510.600(c) of this chapter:
- (1) No. 000859 for use of the product described in paragraph (a)(1) of this section, as in paragraph (c)(1) of this section; and for use of the product described in paragraph (a)(2) of this section, as in paragraph (c)(2) of this section
- (2) No. 000859 for use of the product described in paragraph (a)(1) of this section, as in paragraph (c)(1) of this section.
- (c) Conditions of use—(1) Dogs—(1) Amount. 5 pounds (lb) and under,  $\frac{1}{2}$  tablet (17 mg); 6 to 10 lb, 1 tablet (34 mg); 11 to 15 lb,  $\frac{1}{2}$  tablets (51 mg); 16 to 30 lb, 2 tablets (68 mg); 31 to 45 lb, 3 tablets (102 mg); 46 to 60 lb, 4 tablets (136 mg); over 60 lb, 5 tablets maximum (170 mg). Administer directly by mouth or crumbled and in feed.
- (ii) Indications for use—(A) For removal of canine cestodes Dipylidium caninum and Taenia pisiformis.
- (B) For removal of the canine cestode *Echinococcus granulosus*, and for removal and control of the canine cestode *Echinococcus multilocularis*.

- (iii) Limitations—(A) If labeled only for use as in paragraph (c)(1)(ii)(A) of this section: Not intended for use in puppies less than 4 weeks of age. Consult your veterinarian before administering tablets to weak or debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.
- (B) If labeled for use as in paragraph (c)(1)(ii)(B) of this section: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (2) Cats—(i) Indications for use. For removal of feline cestodes Dipylidium caninum and Taenia taeniaeformis.
- (ii) *Dosage*. Cats 4 pounds and under, 11.5 mg; 5 to 11 pounds, 23 mg; over 11 pounds, 34.5 mg.
- (iii) Limitations. Administer directly by mouth or crumbled and in feed. Not intended for use in kittens less than 6 weeks of age. For OTC use: Consult your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism.
- [46 FR 60570, Dec. 11, 1981, as amended at 47 FR 26377, June 18, 1982; 55 FR 2234, Jan. 23, 1990; 58 FR 7864, Feb. 10, 1993; 58 FR 42853, Aug. 12, 1993; 68 FR 57351, Oct. 3, 2003; 69 FR 62181, Oct. 25, 2004; 78 FR 17596, Mar. 22, 2013]

## §520.1871 Praziquantel and pyrantel.

- (a) Specifications. (1) Each tablet contains 13.6 milligrams (mg) praziquantel and 54.3 mg pyrantel base (as pyrantel pamoate), 18.2 mg praziquantel and 72.6 mg pyrantel base (as pyrantel pamoate), or 27.2 mg praziquantel and 108.6 mg pyrantel base (as pyrantel pamoate).
- (2) Each chewable tablet contains 30 mg praziquantel and 30 mg pyrantel pamoate or 114 mg praziquantel and 114 mg pyrantel pamoate.
- (b) Sponsors. See sponsors in §510.600(c) for use as in paragraph (d) of this chapter.
- (1) See No. 000859 for use of tablets described in paragraph (a)(1) of this section for use as in paragraph (d)(1) of this section.
- (2) See No. 051311 for use of tablets described in paragraph (a)(2) of this section for use as in paragraph (d)(2) of this section.
- (c) Special considerations. See §500.25 of this chapter.